




Heidelberg Institute for
Theoretical Studies



Simulations of biomaterials under force

Part I: Collagen


Frauke Gräter, Bogota, 09/2019




MOLECULAR (bio)mechanics – why?




collagen




mineralized tissue




silk




polymer



MOLECULAR (bio)mechanics – why?




MOLECULAR (bio)mechanics – why?



Strain and fracture: force distribution

conventional design tools:
force distribution

in constructions, cars ...




*macroscopic structures:
meters*

Strain and fracture: force distribution

conventional design tools:
force distribution


in constructions, cars ...



*macroscopic structures:
meters*


our method:
force distribution in (bio)molecules at atomistic level

e.g.




*microscopic structures:
~ 10⁻⁹ meters*


in graphene



*W. Stacklies, et al, PLoS Comp Biol, 2009
Costescu et al, BMC Biophys, 2012*



Forces from Molecular Dynamics simulations



1.

$$\begin{aligned}
 E = & \sum_{\text{bonds}} \frac{k_i}{2} (l_i - l_{i,0})^2 \\
 & + \sum_{\text{angles}} \frac{k_i}{2} (\theta_i - \theta_{i,0})^2 \\
 & + \sum_{\text{torsions}} \frac{V_n}{2} (1 + \cos(n\omega - \gamma)) \\
 & + \sum_{i=1}^N \sum_{j=i+1}^N \left(4\epsilon_{ij} \left(\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^6 \right) + \left(\frac{q_i q_j}{4\pi\epsilon_0 r_{ij}} \right) \right)
 \end{aligned}$$

} bonded interactions ↔
} non-bonded interactions ↔


2.

&


$$F_i = \frac{\delta E}{\delta r_i} = m_i \frac{d^2 r_i}{dt^2}$$

Forces from Molecular Dynamics simulations

F'_{ij} force between atom i and j in relaxed state



F_{ij} force between atom i and j in stretched state



change in pairwise forces


$$\Delta F_{ij} = F_{ij} - F'_{ij}$$




*compare: Newton's cradle
(from wikipedia)*

Forces from Molecular Dynamics simulations

Force Distribution Analysis
(Gromacs patch, on github)




change in pairwise forces
 $\Delta F_{ij} = F_{ij} - F'_{ij}$




- with many applications to proteins
- here: internal strain in a structural component of silk fibers

Compare:
force distribution
in large structures




compare: Admal/Tadmor 2010 (LAMMPS), Vanegas/Arroyo 2014 (Gromacs)

MOLECULAR (bio)mechanics – why?




Molecular
Dynamics &
Monte Carlo &
experiments



Collagen:
most abundant protein and
major load-bearing
structure of the body

*Anatomic plate from Laskowski's
"Anatomie normale du corps
humain" (1894), illustrations by
Sigismond Balicki*


How does collagen behave under force?




highly elastic and resistant
against mechanical load

human achilles tendon:
bears 90 MPa (several
times the body weight)

Collagen: let's pull a nano-meter sized part of a fibril



Wanted:
atomistic model
of collagen



Collagen: integrative modelling of all-atom fibril

Agnieszka
Obarska-Kosinska




X-ray structures,
high resolution, 1.2-1.8 Å



Kramer et al. J Mol Biol. 1998 etc.


THEBuScr
triple-helical
builder

JK Rainey, MC Goh



Collagen: integrative modelling of all-atom fibril


X-ray structures,
high resolution, 1.2-1.8 Å



Kramer et al. J Mol Biol. 1998 etc.

THEBuScr
triple-helical
builder

JK Rainey, MC Goh




X-ray fiber diffraction structure, low resolution: 5.2 Å axial/11 Å equatorial



Orgel et al, PNAS 2006

Collagen: integrative modelling of all-atom fibril


X-ray structures,
high resolution, 1.2-1.8 Å



Kramer et al. J Mol Biol. 1998 etc.

THEBuScr
triple-helical
builder

JK Rainey, MC Goh



low resolution template


X-ray fiber diffraction structure, low resolution: 5.2 Å axial/11 Å equatorial



Orgel et al, PNAS 2006

Collagen: integrative modelling of all-atom fibril



X-ray structures,
high resolution, 1.2-1.8 Å



Kramer et al. J Mol Biol. 1998 etc.

THEBuScr
triple-helical
builder


JK Rainey, MC Goh



low resolution template

refined model using distance-restraints from high-res data


X-ray fiber diffraction structure, low resolution: 5.2 Å axial/11 Å equatorial



Orgel et al, PNAS 2006

Collagen: integrative modelling of all-atom fibril



X-ray structures,
high resolution, 1.2-1.8 Å



Kramer et al. J Mol Biol. 1998 etc.

THEBuScr
triple-helical
builder

JK Rainey, MC Goh



refined model integrating low and high resolution data

X-ray fiber diffraction structure, low resolution: 5.2 Å axial/11 Å equatorial



Orgel et al, PNAS 2006


Collagen: integrative modelling of all-atom fibril

X-ray structures,
high resolution, 1.2-1.8 Å




Kramer et al. J Mol Biol. 1998 etc.



Mass spec data



eg Eyre et al, Methods 2008



THEBuScr
triple-helical
builder
JK Rainey, MC Goh



X-ray fiber diffraction structure, low resolution: 5.2 Å axial/11 Å equatorial



Orgel et al, PNAS 2006

Collagen: integrative modelling of all-atom fibril


X-ray structures
high resolution



Kramer et al. J Mol Biol

THEBUS
triple-helix
builder


JK Rainey, MC G



X-ray fiber di

Orgel et al, PNAS 2006

Mass spec data





Up and running (hopefully) soon:
web service to generate collagen fibril
models at wish

Obarska-Kosinska et al, in preparation


ion data

Reactive Molecular Dynamics: spontaneous chemical events on-the-fly

Benedikt
Rennekamp



Molecular Dynamics simulations: bonds can not rupture!



1.

$$\begin{aligned}
 E = & \sum_{\text{bonds}} \frac{k_i}{2} (l_i - l_{i,0})^2 \\
 & + \sum_{\text{angles}} \frac{k_i}{2} (\theta_i - \theta_{i,0})^2 \\
 & + \sum_{\text{torsions}} \frac{V_n}{2} (1 + \cos(n\omega - \gamma)) \\
 & + \sum_{i=1}^N \sum_{j=i+1}^N \left(4\epsilon_{ij} \left(\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^6 \right) + \left(\frac{q_i q_j}{4\pi\epsilon_0 r_{ij}} \right) \right)
 \end{aligned}$$

} bonded interactions \longleftrightarrow
 \longleftrightarrow
 } non-bonded interactions

2.

&

$$F_i = \frac{\delta E}{\delta r_i} = m_i \frac{d^2 r_i}{dt^2}$$

“Reactive” Molecular Dynamics: spontaneous chemical events on-the-fly


Computational scheme:

Kinetic Monte Carlo/Molecular Dynamics KIMMDY

1. Run n MD steps
2. Check forces, calculate rates
3. Do kinetic Monte Carlo step
4. Perform bond scission
5. Go back to 1.

Currently being extended to
electron transfer and other redox-
chemistries

Using **PLUMED/GROMACS**



Rennekamp, Gräter, in review

Monte Carlo methods

stochastic simulation methods, **sample randomly** to get obtain an approximate numerical result for a complex (high-dimensional?) problem

flavors:

- Monte Carlo integration
- Metropolis Monte Carlo
- Quantum Monte Carlo
- kinetic Monte Carlo

applications:

e.g. optimization, simulations of solids/molecules/polymers, weather, astro, finance, law,....



Monte Carlo methods: some history



some history: the name was suggested by Metropolis, at Los Alamos National Laboratories for a method put forward by Fermi/Ulam for calculations of neutron diffusion

Monte Carlo methods: some history



Strength of random sampling:

Calculation of π by Laplace
(based on the Buffon Experiment)

Suppose we have a floor made of parallel strips of wood, each the same width, and we drop a needle onto the floor. What is the probability that the needle will lie across a line between two strips?


*n needles,
h of them crossing the strips
l length of needles
t width of strips*

$$\pi \approx \frac{2l \cdot n}{th}$$



wikipedia

Kinetic Monte Carlo



- Calculate different rates Γ_i
- Draw random number R between 0 and Γ_{tot}
- Carry out corresponding event
- Draw another random number T between 0 and Γ_{tot}
- Update time with $\Delta t = \log(1/T)$

Advantages:

- any time scale reachable (lower rates just lead to higher time steps)

Disadvantages:

- jump in time: no continuous trajectory
- requires good guesses for rates

“Reactive” Molecular Dynamics: spontaneous chemical events on-the-fly


Computational scheme:

Kinetic Monte Carlo/Molecular Dynamics KIMMDY

1. Run n MD steps
2. Check forces, calculate rates
- 3. Do kinetic Monte Carlo step**
4. Perform bond scission
5. Go back to 1.



Currently being extended to
electron transfer and other redox-
chemistries

Using **PLUMED/GROMACS**



Rennekamp, Gräter, in review

Model: How to calculate rupture rates?



Using Bell-Evans model for the potential and Arrhenius kinetics to calculate rupture rates; taking parameters for the potential from force field.


$$V(x) = D[1 - \exp(\beta(x - x_0))]^2$$

$$V_{eff} = V(x) - x \cdot V'(x_{curr})$$

$$\Delta E = V_{eff}(x_{max}) - V_{eff}(x_{min})$$


$$\Rightarrow k_{rupture} = A \cdot \exp(-\Delta E/k_B T)$$

Weakest bonds and their rates



$$E_{\text{dis}}(\text{CA-N}) = 348 = E_{\text{dis}}(\text{CA-C})$$

It works: Rupture preferentially occurs around crosslinks





Molecular Biomechanics

Ana Herrera-Rodriguez
Nicholas Maragakis
Florian Franz
Fabian Kutzki
Christopher Zapp
Agnieszka Obarska-Kosinski
Fan Jin
Benedikt Rennekamp
Isabel Martin
Anna Schroeder

\$\$:
Klaus Tschira foundation
DFG, Volkswagen Foundation,
AvH,
Toyota

Collagen: integrative modelling of atomistic system


X-ray structures,
high resolution, 1.2-1.8 Å
1Z1R, 1A3J, 1CAG, 1QSU



Kramer et al. J Mol Biol. 1998
Okuyama, Biopolymers. 2009
etc.

THEBuScr
triple-helical
builder

JK Rainey, MC Goh



X-ray fiber diffraction structure 3HR2
low resolution: 5.2 Å axial/11 Å equatorial
Orgel et al, PNAS 2006

